

1. The function $f(x) = \begin{cases} ax^2 + bx & \text{if } x \leq 1 \\ x + a^2 & \text{if } x > 1 \end{cases}$

is twice differentiable everywhere. Then $a^2 + b^2 =$

- a) 1
- b) 0
- c) $\frac{5}{4}$
- d) 2
- e) 5

2. If $f(x) = (2x - 1)^{\frac{2}{3}}$, then the equation of the vertical tangent to the graph of f is

- a) $x = \frac{1}{2}$
- b) $x = -\frac{1}{2}$
- c) $x = \frac{2}{3}$
- d) $x = -\frac{2}{3}$
- e) $x = \frac{4}{3}$

3. The equations of the horizontal tangents to the curve $y = x^3 - 3x - 2$ are
- a) $y = 0$ and $y = -4$
 - b) $y = 1$ and $y = -1$
 - c) $x = 1$ and $x = -1$
 - d) $y = -4$ and $y = 1$
 - e) $y = 0$ and $y = -1$
4. At how many real values of x does the curve $y = x^6 - 3x^2 + x + 5$ have a tangent line parallel to the line $y = x$?
- a) 3
 - b) 1
 - c) 2
 - d) 4
 - e) 5

5. If the tangent line to the graph of $f(x) = \frac{2x}{2x+1}$ at the point (α, β) is $y = 2x + 1$, then $\beta^2 =$

- a) 1
- b) 2
- c) 3
- d) 4
- e) 5

6. If $f(x) = xe^x$ and n is a positive integer, then $f^{(n)}(1) =$

- a) $(n+1)e$
- b) ne
- c) $(n-1)e$
- d) $(n+2)e$
- e) $ne+1$

7. If $y = \frac{1 + \sin x}{1 + \cos x}$, then $\frac{dy}{dx} =$

a) $\frac{1 + \sin x + \cos x}{(1 + \cos x)^2}$

b) $\frac{\sin x + \cos x}{1 + \cos x}$

c) $\frac{\sin x + \cos x}{(1 + \cos x)^2}$

d) $\frac{1 + \sin x}{(1 + \cos x)^2}$

e) $\frac{2}{1 + \cos x}$

8. $\lim_{\theta \rightarrow 1} \frac{\sin(\theta - 1)}{\theta^2 + \theta - 2} =$

a) $\frac{1}{3}$

b) 0

c) $\frac{1}{2}$

d) 2

e) 1

9. If $y = \sin(x^2)$ and $x = \cos t$, then $\frac{dy}{dt} =$

- a) $-\sin 2t \cos(\cos^2 t)$
- b) $\sin 2t \cos(\cos^2 t)$
- c) $-\sin t \cos(\cos^2 t)$
- d) $-\sin 2t \cos^3 t$
- e) $\sin 2t \cos^3 t$

10. Let f and g be differentiable functions and $h(x) = f(x^2g(x))$. If $g(2) = -2$ and $g'(2) = 2$, then $h'(2) =$

- a) 0
- b) -2
- c) 2
- d) 3
- e) -3

11. The equation of the tangent line to the curve given implicitly by

$$\sqrt{x+y} = y^2$$

at the point $(0, 1)$ is

- a) $3y - x = 3$
- b) $2y + x = 1$
- c) $3y + x = 3$
- d) $2y - x = 2$
- e) $2y + x = 3$

12. The equation of the normal line to the curve $y = \tan^{-1}(\sqrt{x-1})$ at $x = 2$ is

- a) $y = -4x + 8 + \frac{\pi}{4}$
- b) $y = \frac{1}{4}x - \frac{1}{2} + \frac{\pi}{4}$
- c) $y = 4x - 8 + \frac{\pi}{4}$
- d) $y = -\frac{1}{4}x + \frac{1}{2} + \frac{\pi}{4}$
- e) $y = -4x + 8 - \frac{\pi}{4}$

13. If $f(x) = (x^2 + 2x)^{50}$, then $f^{(100)}(1) =$

- a) $100!$
- b) 100
- c) 0
- d) $3(99!)$
- e) $2(50!)$

14. The slope of the tangent line to the graph of $y = (2x + 1)^{\sin 3x}$ at $x = \frac{\pi}{6}$ is

- a) 2
- b) $4\left(\frac{\pi}{3} + 1\right)$
- c) 6
- d) $2\left(\frac{\pi}{3} + 1\right)$
- e) $\frac{4}{\frac{\pi}{3} + 1}$

15. If $y = \frac{(x+2)^2(2x-1)^3}{\sqrt{x+1}}$, then $y'(0) =$

- a) 22
- b) $-\frac{11}{2}$
- c) 44
- d) 24
- e) -11

16. The position function of a particle moving along a line is

$$s(t) = \sin t + \cos t$$

where t is measured in seconds and s in meters. The total distance traveled by the particle in the interval $[0, \pi]$ is

- a) $2\sqrt{2}$ meters
- b) 2 meters
- c) 4 meters
- d) $2\sqrt{2} + 2$ meters
- e) $2\sqrt{2} - 2$ meters

17. The position function of a particle moving along a line is

$$s(t) = t^3 - 6t^2 + 9t \quad (0 \leq t \leq 5).$$

The time interval(s) where the particle is moving forward is (are)

- a) (0, 1) and (3, 5)
 - b) (0, 3)
 - c) (0, 3) and (4, 5)
 - d) (1, 3)
 - e) (0, 2) and (3, 5)
18. The two equal sides of an isosceles triangle have length $4m$. If the angle between them is increasing at a rate of 0.06 rad/s , then the rate at which the area of the triangle is changing when the angle between the sides of the triangle is $\frac{\pi}{3}$ equals
- a) $0.24 \text{ m}^2/\text{s}$
 - b) $-0.24 \text{ m}^2/\text{s}$
 - c) $2.4 \text{ m}^2/\text{s}$
 - d) $-2.4 \text{ m}^2/\text{s}$
 - e) $0.024 \text{ m}^2/\text{s}$

19. If a snow ball melts so that its surface area decreases at a rate of $1 \text{ cm}^2/\text{min}$, then the rate at which the diameter changes, when the diameter is 10 cm equals

Hint: Surface area of a sphere = $4\pi r^2$

- a) $\frac{-1}{20\pi} \text{ cm/min}$
- b) $\frac{1}{20\pi} \text{ cm/min}$
- c) $\frac{-1}{40\pi} \text{ cm/min}$
- d) $\frac{1}{40\pi} \text{ cm/min}$
- e) $\frac{-1}{10\pi} \text{ cm/min}$

20. The equation of the tangent line to the graph of $y = \ln x$ and passes through the origin is

- a) $e y = x$
- b) $y = e x$
- c) $y = \frac{1}{e}(x - 1)$
- d) $y = \frac{1}{e}(x + 1)$
- e) $y = 2 e x$