



1. Using **the method of cylindrical shells**, set up, but do not evaluate, an integral for the volume of the solid obtained by rotating

- (a) [**7 points**] the region bounded by the curves  $y = \sqrt{x}$ ,  $y = x - 2$  and  $y = 0$  about the  $x$ -axis. [Sketch the region and a typical rectangle].



- (b) [**7 points**] the region bounded by the circle  $x^2 + y^2 = 1$  about the line  $x = -1$ . [Sketch the region and a typical rectangle].



2. [**5 points**] Find the average value of the function  $f(t) = \tan t \sec t$  over the interval  $\left[0, \frac{\pi}{4}\right]$ .

3. Determine whether the integral is convergent or divergent. If it is convergent, find its value.

(a) [6 points]  $\int_0^9 \frac{1}{x\sqrt{x}} dx$ .

(b) [8 points]  $\int_0^{+\infty} x e^{-10x} dx$ .

4. [6 points] Determine whether the sequence  $\left\{ \frac{(-1)^n \sqrt{n}}{n+7} \right\}_{n=1}^{+\infty}$  is convergent or divergent. If it is convergent, find its limit.

5. [7 points] Use geometric series to write the number

$$1.2\overline{13} = 1.2131313\dots$$

as a ratio of two integers.

6. Evaluate the following integrals:

(a) [**9 points**]  $\int x(\ln x)^2 dx.$

(b) [**10 points**]  $\int \frac{x^3}{\sqrt{4-x^2}} dx.$

(c) [12 points]  $\int \frac{x^3 + 1}{x^3 + x} dx.$

(d) [10 points]  $\int \frac{\sec x}{2 + \tan x} dx.$  Hint: Use the substitution  $t = \tan\left(\frac{x}{2}\right).$

7. Determine whether the series is convergent or divergent. If it is convergent, find its sum.

(a) [6 points]  $\sum_{n=1}^{+\infty} \left(\frac{1}{2}\right)^{\frac{1}{n^2}}$  .

(b) [7 points]  $\sum_{n=1}^{+\infty} [\tan^{-1}(2n-1) - \tan^{-1}(2n+1)]$  .